

Progress Report Narrative

Recipient Name: Dr. Michael J. Weber, Iowa State University

Agreement Number: F16AP00791

Project Title: Asian carp population ecology in tributaries of the Upper Mississippi River

Report Type: Interim

Period Covered: Aug 1, 2016 – July 31, 2017

Project Narrative: (documents progress made in achieving the objectives of project work plan(s) approved in application)

Provide the following information:

1. Comparison of actual accomplishments (outputs, outcomes) with the anticipated outputs/outcomes.

The Goals and Objectives of this project are to:

1. Evaluate adult population characteristic (abundance, distribution, size structure, condition) and dynamics (recruitment, growth, mortality) of Asian carp in pools 14, 15, 16, 17, 18, 19, and 20 of the Upper Mississippi River.

2. Evaluate Asian carp reproduction in pools 14, 15, 16, 17, 18, 19, 20 and the contribution of the Wapsipinicon, Rock, Iowa, Skunk, and Des Moines rivers to Asian carp reproduction (egg, larval and juvenile densities).

We have conducted all field sampling for this project and are on target to complete sample processing and data analysis by the completion date of April 1, 2018 as stated in amendment 1 of F16AP00791. Egg and larval ichthyoplankton samples were collected throughout the summer in pools 14-20 to monitor Asian Carp reproduction across the leading edge of the invasion. Adult Asian Carp were sampled during fall 2016 in pools 14-20 to document variation in population characteristics across the leading edge of the invasion. Specific details and preliminary results of our actual accomplishments to date are provided in the report (see below).

2. Reasons why anticipated outcomes were exceeded or not met.

NA – all project goals and objectives will be met.

3. If applicable, problems encountered during the performance period, which may interfere with meeting program/project objectives. (List N/A if no problem exists.)

Sample processing has taken longer than anticipated due to the large volume of chlorophyll, zooplankton, Ichthyoplankton (>700 samples/each), and adult samples that were collected and large number of zooplankton, egg, and larval fish present in the samples. Despite delays, all anticipated outcomes will still be met. We were also not able to sample on a few dates and a few sites due to environmental conditions (e.g., water levels too low or high for safe sampling) and mechanical issues. These instances were few and noted in the report.

4. List proposed remedies if problem(s) exist (s) as indicated in item 3. [List N/A if not applicable]

I am in contact with the Limnology Lab at Iowa State University to gauge their interest in processing the chlorophyll and zooplankton samples and at what associated cost. If they are able to process these samples for us, we will have more resources to focus on process the fish samples in a more timely manner.

5. Information on the rate of expenditure versus progress on project.

Cost for sample processing have been higher than anticipated due to the higher egg and larval densities and debris in the ichthyoplankton samples than anticipated. As the original project completion date was August 1, 2017, project funds have nearly been exhausted.

6. If applicable, information on equipment purchased during the reporting period. [List N/A if no equipment purchased during reporting period]

NA - No equipment has been purchased for the project.

7. Any additional pertinent information, including, when appropriate, analysis and information of cost overruns or high unit costs or unanticipated economics. [List N/A if not applicable]

NA

Annual Progress Report to:
United States Fish and Wildlife Service

Distribution and Population Dynamics of Asian Carp in Iowa Rivers

Submitted by

Aaron J. Matthews, Michael J. Weber, and Clay L. Pierce

Department of Natural Resource Ecology and Management
Iowa State University

October 2017

Table of Contents

ABSTRACT	3
INTRODUCTION	4
BACKGROUND	4
RESEARCH NEEDS.....	6
OBJECTIVES.....	6
STUDY AREA	6
METHODS	7
EGG AND LARVAL SAMPLING.....	7
ADULT SAMPLING	8
ENVIRONMENTAL CONDITION DATA COLLECTION.....	8
PRELIMINARY RESULTS	9
ASIAN CARP REPRODUCTION.....	9
ADULT ASIAN CARP POPULATION CHARACTERISTICS AND DYNAMICS	10
ENVIRONMENTAL CONDITION DATA	10
PRODUCTS RESULTING FROM THIS PROJECT	11
LITERATURE CITED	12
TABLES AND FIGURES	16
<i>Table 1. Eggs of 13 fish species captured in 2016 by date and river. River abbreviations present for a species on a sampling date denotes genetic confirmation of that species collected at that location on a sampling date. W= Wapsipinicon River, R= Rock River, I= Iowa River, S= Skunk River, D= Des Moines River, U= Upper Mississippi River.</i>	<i>16</i>
<i>Table 2. Electrofishing catch per unit effort (CPUE; mean ± 1 SE) of Silver (SC) and Bighead Carp (BC) by location and date captured during 2016.....</i>	<i>17</i>
<i>Figure 1. Sampling sites (diamond) and lock and dam locations (bold bar) along the Mississippi River and major tributaries.</i>	<i>18</i>
<i>Figure 2. Density (mean ± 1 SE) of eggs (top) and age-0 fish (bottom) during 10 day sampling sessions.....</i>	<i>19</i>
<i>Figure 3. Densities (mean ± 1 SE) of eggs (top) and age-0 fish (bottom) from each site during 2016.</i>	<i>20</i>
<i>Figure 4. Egg densities (mean ± 1 SE) captured from backwater, channel border, and thalweg habitats during 2016.</i>	<i>21</i>
<i>Figure 5. Age-0 fish density (mean ± 1 SE) in upstream (black), tributary (dark grey, forward slash), and downstream (light grey) locations associated with the Des Moines (left), Skunk (middle), and Iowa (right) river confluences. Upstream = 1 km upstream of confluence in the Mississippi River, Tributary = 1 km upstream of confluence in the tributary, and Downstream= 1 km downstream of confluence in the Mississippi River.</i>	<i>22</i>
<i>Figure 6. Silver Carp length-frequency distributions and proportional size distribution (PSD; P- Preferred, M- Memorable, T-Trophy) indices of fish collected during 2016 from the Des Moines River at Cliffland (CLF) and at the mouth (MTH) in addition to Iowa River and Skunk.....</i>	<i>23</i>
<i>Figure 8. Chlorophyll a measurements (mean ± 1 SE) of each site.....</i>	<i>24</i>

Abstract

Invasive Silver Carp (*Hypophthalmichthys molitrix*) and Bighead Carp (*H. nobilis*; collectively called Asian Carp) are expanding throughout the Upper Mississippi River Basin (UMRB) and are of great concern due to their potential economic and ecological impacts. Pooled sections on the Upper Mississippi River associated with lock and dams may be poor habitats for reproduction and recruitment due to their lentic characteristics and perceived lack of adequate spawning habitat compared to more free-flowing unimpounded sections in the lower Mississippi River where evidence of reproduction has been documented. However, Iowa interior rivers connected to pooled sections of the Upper Mississippi River possess several requirements needed for successful spawning and observations of adults are becoming more prevalent. Unfortunately, little is known regarding the basic ecology and reproductive status of Asian Carp populations in these tributary systems. In order to properly make management decisions, information on reproductive success and factors that regulate populations must be understood. In this study, we evaluate 1) reproduction patterns and 2) adult population characteristics and dynamics of Asian Carp in the Mississippi, Des Moines, Skunk, Iowa, Rock, and Wapsipinicon rivers in southeastern Iowa. In 2016, 744 ichthyoplankton tows captured 26,240 eggs and 42,068 age-0 fishes from the Wapsipinicon, Rock, Iowa, Skunk, Des Moines and Mississippi rivers. Age-0 fish are currently being identified to determine densities of Asian Carp versus other fishes. The largest number of eggs collected in a sampling event occurred on May 29th and 30th 2016 when 22,689 eggs (86% of all eggs collected during 2016) were collected. The majority of eggs during this sampling event were collected from tributaries (17,509 of 22,689 eggs, 86%) compared to the Upper Mississippi River. Among tributaries, the majority of eggs were collected from the Skunk River mouth (15,553 eggs). In contrast, the majority of age-0 fish (39,700 of 42,068, 96%) were captured from the Mississippi River in 2016. Egg densities peaked in late May while age-0 fish densities were greatest during late June. Genetic analysis verified that Asian Carp eggs were captured in the Upper Mississippi River as far north as Pool 18 in addition to the Iowa and Skunk river mouths. Larval sampling for the 2017 field season is complete but samples are still being processed. Egg, larval, and adult samples will continue to be processed and additional data analysis will be conducted in 2017 and 2018 to evaluate spatiotemporal patterns of reproduction and adult population characteristics. In addition to eggs and age-0 fish, a total of 186 adult Asian Carp were collected from the Mississippi, Des Moines, Skunk, Iowa, and Rock rivers during fall 2016. The majority of these fish were Silver Carp (96%) and most were collected in Pool 20 at the mouth of the Des Moines River (94%). Four Bighead Carp were collected throughout the UMRB during 2016, one at Cliffland on the Des Moines and three at the Des Moines River Confluence. Additionally, two Grass Carp were collected at Cliffland on the Des Moines River. Additional adult sampling will occur during fall 2017.

Introduction

Background

Ecological communities worldwide are becoming more uniform through the introduction and subsequent establishment of non-native species into novel areas through anthropogenic activities (Rahel 2002). Intentional introductions commonly occur to provide societal benefits such as food, recreation, and biological control (Pimentel et al. 2000). Additionally, advancements in transportation and worldwide commerce have increased unintentional introductions (Rahel 2002). In the United States, approximately 50,000 non-native species introductions have occurred with varying success and impacts (Pimentel et al. 2005, Sagoff 2005). Nevertheless, a single non-native species can alter ecosystem structure and function and have costly economic consequences (Macisaac 1996, Pimentel et al. 2005, Weber and Brown 2009). Economic losses due to non-native introductions are estimated at US\$120 billion a year; however, actual costs are likely much higher because monetary costs associated from species extinctions, loss of ecosystem services, and aesthetic values are not easily assessed (Pimentel et al. 2005). Likewise, ecological costs may be much greater than economic costs but are difficult to quantify and assess because of lag times between invasion and empirically confirmed impacts to the environment (Gido and Brown 1999, Stohlgren and Schnase 2006).

Non-native fishes are one of the most introduced groups of aquatic animals that have resulted in negative ecological effects in riverine ecosystems (Gozlan 2008). Rivers naturally provide an invasion highway for fishes to expand from the point of introduction. Furthermore, modifications to rivers for navigation have connected previously separate waterways, facilitating inter-basin movement and the spread of invaders into additional novel habitats (Leuven et al. 2009). For example, the Mississippi River Basin covers roughly 40% of the lower 48 U.S. states with thousands of river miles (USACE 2011) and connects to the previously separated Great Lakes basin via the Chicago Area Waterway System, the Ohio-Erie Canal and other man-made structures (USACE 2014).

At least 83 non-native fishes have become established in the Upper Mississippi River Basin (UMRB) as a result of dispersal from other basins or by direct introduction from anthropogenic activities (Rasmussen 2002). Two of the more recent and widely recognized invaders to the UMRB are Silver Carp (*Hypophthalmichthys molotrix*) and Bighead Carp (*H. nobilis*; collectively referred to as Asian Carp). These species have become abundant and threaten the integrity of the UMRB and any connected aquatic ecosystems (Irons et al. 2009). Asian Carp were imported during the 1970s into the United States for food consumption and biological control in aquaculture facilities (Freeze and Henderson 1982). In the 1970s, individuals thought to have escaped during flooding events were observed in several rivers within Arkansas. Due to their high reproductive capabilities and long distance migrations (DeGrandchamp et al. 2008), these fish quickly became established and now inhabit more than 20 states throughout the Mississippi, Missouri, Ohio, and Illinois river basins (Kolar et al. 2007, Baerwaldt et al. 2013, Deters et al. 2013). By the mid-1980s, Asian Carp were caught in the pooled sections of the UMRB (Kolar et al. 2007) with the first observations of Asian Carp in Iowa occurring in 1986 when Silver Carp were captured in the UMRB below lock and dam 19 (LD19)

near Keokuk (Irons et al. 2009). A year later, Bighead Carp were captured near the mouth of Yellow Springs Creek north of Burlington, IA (Irons et al. 2009). Since the initial observations in Iowa, Asian Carp adults have been sighted in several additional UMRB tributaries in Iowa such as the Des Moines, Skunk, Iowa, and Cedar rivers (Bruce 1990, United Press International 2011, Irons 2012, Camacho 2016, Sullivan 2016).

Currently, southeastern Iowa appears to be on the leading edge of Asian Carp expansion in the UMRB. Substantially higher adult catch rates of both Silver and Bighead Carp occur below LD19 than above, suggesting this structure and other lock and dams on the Mississippi River may serve as a partial migration barriers (Wilcox et al. 2004). For example, the UMRB water level is regulated at each dam in order to maintain a navigation channel by reducing or eliminating the amount of water discharged, leaving passage through the locks as the only means of fish movement during low river discharge periods. However, dam gates are lifted during higher discharge events, that facilitate fish passage (Garvey et al. 2010). It is also during these high discharge events that Asian Carp exhibit some of their highest movement rates, especially during annual spring runoff and associated peak discharge events when temperatures are below or within the spawning optimum, suggesting movement may be associated with spawning migration behavior (Jennings 1988, Peters et al. 2006, DeGrandchamp et al. 2008). Furthermore, Asian Carp can quickly make long distance migrations (DeGrandchamp et al. 2008) indicating that these fish are capable of dispersal into new locations.

Although Asian Carp may be able to navigate lock and dams on the UMRB, pooled sections between these structures may provide unsuitable spawning habitats for these species. Asian Carp are highly fecund (up to 3.5 million eggs per female; Garvey et al. 2006) and have short gestation periods (Chapman and George 2011). Thus, only a few adult individuals may be needed to quickly establish an abundant population (Crawley et al. 1986). Despite adult Asian Carp being detected above LD19 up to St. Paul, MN, USA, their populations have remained low, suggesting reproduction may be limited in these reaches. Pooled sections associated with lock and dams exhibit reservoir-like characteristics that are more lentic in nature resulting in lower Asian carp reproduction than in unregulated sections where lock and dams are absent (Lohmeyer and Garvey 2009). In contrast, established Asian Carp populations in tributary systems, such as the Illinois River, can have high recruitment and adult populations have increased exponentially in abundance within a decade (Sass et al. 2010). Yet, Asian Carp abundance in the tributaries of the UMRB are much lower compared to the Illinois River and it is unknown whether or not these systems are suitable for reproduction.

Successful Asian Carp spawning depends on adults finding suitable habitat of sustained, high flow or increasing discharge when water temperatures are between 17 and 30°C (Kolar et al. 2007). Continuous river flow of at least 25 km may be necessary to suspend the semi-buoyant eggs for a 24 h period or until larvae successfully hatch (Krykhtin and Gorbach 1981, George and Chapman 2013, Murphy and Jackson 2013). In most areas of the UMRB, reaches between dams with sufficient sustained velocities of 0.3 to 3.0 m/s and turbulence to keep eggs in suspension do not exist or are poorly suited for egg survival (Lohmeyer and Garvey 2009). However, age-0 Asian Carp have been documented in tributaries such as the Cache River (a tributary to the Ohio River; Burr et al. 1996) and the Illinois River (a tributary to the Mississippi River; DeGrandchamp et al. 2007). Additionally, tributaries are known to be associated with

spawning activity in their native range in the Yangtze River (Yi et al. 1988) and in varying capacities in the Missouri River (Schrank et al. 2001) and Illinois River (DeGrandchamp et al. 2007) where they are introduced. Successful establishment and reproduction in tributaries could provide sources of recruitment for pooled sections of the UMRB and other areas of poor reproduction.

Research Needs

Successful expansion and establishment of Asian Carp populations within the UMRB depends on the ability of adults to find adequate conditions of temperature and long fetches of sustained, high flow. Despite perceived poor conditions for successful Asian Carp reproduction in the Upper Mississippi River, tributaries could provide adequate conditions for reproduction, resulting in population expansion along the leading edge of the invasion. Reproductive success on an invasion front can increase exponentially through time (Chick and Pegg 2001, Sass et al. 2001) until establishment occurs (Hayer et al. 2014). Some systems where Asian Carp are starting to establish can have sporadic reproduction (Irons et al 2011) leading to either a surge or decline in population. Although adult Asian Carp density appears low, successful reproduction of Asian Carp has been established in previous research at Iowa State within the Des Moines, Skunk, and Iowa rivers (Camacho 2016). Further evaluation of factors affecting reproduction and recruitment in Iowa tributaries of the Mississippi River in associated with annual variation in environmental conditions is needed to better understand Asian Carp population dynamics in these systems and potentially develop management strategies for these invasive fishes. By understanding more about factors affecting reproduction and recruitment within the tributaries of the UMRB, potential increases of Asian Carp presence, or newly established residence in UMR could be detected early. Spatial and temporal distribution of Asian Carp eggs and larvae will help to locate spawning habitat, help determine reproductive cues, and provide insight between environmental variables and survival of larvae and juveniles.

Objectives

The objectives of this study are to evaluate Asian Carp reproduction (egg, larval and juvenile densities) and adult population characteristics (abundance, distribution, size structure) in Pools 14-20 of the UMR with a focus on confluences with the Des Moines, Skunk, Iowa, Rock, and Wapsipinicon rivers.

Study Area

The UMR flows 2,320 river kilometers (RKM) from Lake Itasca, Minnesota to the Ohio River confluence near Cairo, Illinois. Legislative approval by US Congress in 1905 and 1935 altered the UMR by forming a series of dams, levees, wing dikes and other structures to control flooding and provide commercial navigation (Garvey et al. 2010). The UMR dams are preceded by a series of pools, starting at Pool 1 formed by Lock and Dam 1 located near St. Paul Minnesota to Pool 26 above Lock and Dam 26 near Alton, Illinois. The flows of the UMR have shifted from historically lotic to lentic in nature (Garvey et al. 2003). Despite the increase in side

channels, islands and backwaters from the system of dams, they inhibit fish movements for reproduction or expanding their range (Southall and Hubert 1984, Zigler et al. 2004). At low flows when the dams close to retain water, the only passage available is through the series of locks when boats move between pools. Disruption of river continuity could affect species such as Asian Carp that require long stretches of connected habitat for one or more life stages as it does in native species. Egg and larval Asian Carp have been documented as far north as Pool 16 (Larson et al. 2017). Anthropogenic alterations also occur within the major tributaries within the Upper Mississippi River.

The Des Moines, Skunk, Iowa, and Wapsipinicon rivers all contain lowhead dams within 90 km of the confluence of the Mississippi River, potentially inhibiting reproduction within the tributaries. There are low head dams greater than 90 km from the confluence with the Mississippi River, and a split dam located within 10 km from the confluence. Catchment areas range between 6,565 km² for the Wapsipinicon River to 37,296 km² for the Des Moines River. The Des Moines River and the Rock River are the only two rivers with basins within multiple states, Iowa and Minnesota, and Illinois and Wisconsin, respectively. Previous sampling at Iowa State University has found egg and larval Asian Carp within the Des Moines, Skunk, and Iowa tributaries and adults in the Des Moines, Skunk, Iowa and Cedar rivers (Camacho 2016).

Methods

Egg and Larval Sampling

Asian Carp eggs and age-0 fish were sampled from 2016 and 2017 at 18 locations (Figure 1) approximately every 10 days depending upon river conditions from the end of April until September 2016 and 2017 (14 sessions during 2016, with 54 tows per session). Sampling was not conducted when water levels were too high for safe boating or too low for boat access. Ichthyoplankton (0.5 m diameter net, 500 µm mesh) tows were conducted at the surface at a constant boat speed relative to the shoreline up to four minutes depending on debris load. A General Oceanics flowmeter (Model 2030R) was mounted in the mouth of the net to estimate volume (m³) of water filtered during each tow. Three tows were conducted at each site parallel to river flow: the first tow was in the main thalweg for drifting eggs and larvae (<24 hours post fertilization), the second tow occurred near channel borders where water velocity is moving downstream slower than the thalweg, and the third was in an adjacent backwater area for mobile larvae (>24 hours post fertilization). After each tow, ichthyoplankton net contents were rinsed toward the cod end, placed in sample jars, and preserved in 95% ethanol.

In the laboratory, eggs and age-0 fish (larvae and juveniles) were separated from debris. Asian Carp larvae are difficult to distinguish among species and are being identified to genus using meristic and morphometric characteristic (Tweb et al. 1990, Chapman 2006, Chapman and George 2011). A portion of eggs (723 eggs) were randomly selected across sampling dates and locations and identified to species using mitochondrial DNA sequencing at University of Wisconsin-Stevens Point. Age-0 fishes were first categorized as larval or juveniles based on fin development. Fish recognized as having a full complement of fins are categorized as juvenile fish. All age-0 fish are being identified to the lowest possible taxa using morphometric and meristic characteristics described in literature (Auer 1982).

Adult Sampling

Sampling for adult Asian Carp occurred September through November in 2016 at 8 sites in the Mississippi, Des Moines, Skunk, Iowa, Rock, and Wapsipinicon rivers (Figure 1) using daytime boat electrofishing. Asian Carp are notoriously difficult to capture; however, electrofishing is more effective than trammel nets at capturing Silver Carp at these sampling locations (Wanner and Klumb 2009; Sullivan et al. 2017). Thus, electrofishing (pulsed DC; amps 4-13, voltage 100-500) was used to target channel border and backwater areas less than 4 m deep where Asian Carp have previously been shown to typically inhabit (DeGrandchamp et al. 2008). Electrofishing transects (varying effort and transect numbers) are conducted until approximately 150 Silver Carp are captured (Pool 20) or until all available habitat at the site is been sampled.

Asian Carp were identified as a Silver, Bighead, or Silver x Bighead Carp hybrid using meristic and morphometric features (Kolar et al. 2007), weighed (0.001 kg), measured (total length; 1 mm), and the first pectoral fin ray on each side and lapilli otoliths (up to 150 fish/site) were removed for age and growth analysis. Sex is determined based on visual inspection of gonads (male, female, immature, or unknown).

Lapillus otoliths were air dried at room temperature for at least four weeks following collection before being mounted in epoxy. A 1-mm thick cross section at the nucleus was cut using a Buehler Isomet low-speed saw (Isomet Corporation, Springfield, VA) with the anterior portion of the otolith oriented perpendicular to the blade. Wetted 2,000-grit sandpaper was used to polish each side of the cross section. The section was then placed in immersion oil to improve clarity and annuli viewed under a dissecting microscope with transmitted light. Lapillus otoliths were independently aged by two experienced readers with no knowledge of fish length, estimated age of other structure, or source river. If the readers disagreed, then a common age was decided in unison.

Environmental Condition Data Collection

Water temperature, discharge, and stage height were obtained from field measurements and gauging stations on each river. Discharge (m^3/s) and stage height were obtained from U.S. Geological Survey gauging stations across Iowa near Tracy, Ottumwa, Keosauqua, Augusta, Merrimac, Wapello, Iowa City, and Marengo. Water temperature (Yellow Springs Instruments 550A) and conductivity (EC400 ExStik 2 Conductivity Meter) were collected at each site in the thalweg during each sampling period. Across the UMR, river flow data were obtained from various United States Geological Survey (USGS), Advanced Hydrologic Prediction Service (APHS), and National Oceanic and Atmospheric Administration (NOAA) monitoring and gaging stations closest to sampling sites.

Chlorophyll *a* and zooplankton were collected in conjunction with each Asian Carp ichthyoplankton sampling event. Samples were collected from a stationary boat position in thalweg, side channel and backwater habitats. Triplicate zooplankton samples were collected at each site with an integrated tube sampler (5 cm diameter, 50 cm length), filtered through a 63- μm mesh sieve, combined into a composite sample, and preserved using Lugol's solution. Chlorophyll *a* was measured by filtering approximately 100 mL of water through a GF/F

Whatman® glass fiber filter (47-µm porosity) that were placed on ice in the field and frozen in the laboratory. In the laboratory, zooplankton samples were identified to suborder or family and enumerated for total density (number/L). Chlorophyll *a* was extracted with 90% acetone and quantified using an Trilogy Laboratory Fluorometer (Turner Designs) to obtain chlorophyll concentrations (µg/L).

Preliminary Results

Asian Carp Reproduction

In 2016, a total of 744 ichthyoplankton tows were completed. Eggs were collected during every sampling session cumulating in a total of 26,240 eggs being collected across dates and sites in 2016. The largest number of eggs collected in a sampling event was during May 29th and 30th when a total of 22,689 eggs, 86% of the eggs collected in 2016, were captured (Figure 2). Eggs were collected in each river and every site except the backwater habitat of the Wapsipinicon mouth. A total of 495 tows were taken from the Mississippi River that resulted in a total of 8,731 eggs. An additional 84 tows were taken within the tributary mouths that captured 17,509 eggs: 14 eggs were captured in the Wapsipinicon River confluence, 174 eggs were captured in the Rock River confluence, 1,496 eggs were captured in the Iowa River confluence, 15,533 eggs were captured in the Skunk River confluence, 197 eggs were captured in the Des Moines River confluence, and 95 eggs were captured in the Des Moines River at Keosauqua. Mean egg density by site was highest within the Skunk River (SKK-MTH) and lowest at the mouth of the Wapsipinicon River (Figure 3). Within a site, similar egg densities were captured within the backwater, channel boarder, and thalweg habitats (Figure 4).

Genetic results from eggs collected during 2016 confirm the reproduction of 14 different fish species (Table 1). Asian Carp spawning was not directly observed in the field, but egg genetics confirmed Asian Carp eggs were collected on May 29th. Bighead and Silver Carp eggs were identified during May 29th within the Iowa and Skunk river confluences in addition to the Mississippi River downstream of the Skunk River. On May 29th, Silver Carp eggs were found in an additional location within the Mississippi River downstream of the Iowa River. Grass Carp eggs were found intermittently from May 29th to September 9th within, upstream, and downstream of the Iowa and Skunk river confluences. Grass Carp eggs identified on August 18th were only found within the Upper Mississippi River within Pool 17. Grass Carp eggs identified on August 28th were found within the Mississippi River upstream and downstream of the Wapsipinicon, Rock, Iowa, and Skunk River confluences and within Pools 15 and 17.

A total of 42,068 age-0 fish (combination of larvae and juveniles) were captured with ichthyoplankton tows throughout 2016. The highest densities of age-0 fish were collected on June 18 (5.52 ± 2.33 SE, 6,828 fish) and 28 (5.30 ± 1.06 SE, 7,167 fish; Figure 2). The two sessions with the highest mean density of age-0 fish were captured within thirty days of the highest eggs density in 2016. The lowest density of all the sessions (0.46 ± 0.29 SE) was captured during August 28th.

Age-0 fish were sampled from every river during 2016. In contrast to eggs, the majority of age-0 fish (39,700, 94%) were collected from sites within the Mississippi River, whereas only 2,368 (5%) were collected from within tributaries. However, with the exception of the

Wapsipinicon River, downstream tributary sites had higher age-0 fish densities than upstream sites, suggesting a large number of age-0 fish may have originated from tributaries (Figure 3). The highest density of age-0 fish within the Mississippi River (6.88 ± 3.27 SE) were collected downstream of the Des Moines River (UMR-DNW) whereas the lowest densities were collected from WAP-MTH (0.16 ± 0.05 SE). Despite age-0 fish being predominantly captured in backwater habitats during 2014-2015, age-0 fish densities were highest in backwater sites at only 5 of the 18 sites during 2016 whereas densities were highest in the thalweg at 6 of the 18 sites and in channel borders at 5 of the 18 sites (Figure 5).

Adult Asian Carp Population Characteristics and Dynamics

A total of 186 Asian Carp were collected from the Des Moines, Skunk, Iowa, and Rock River sites during a total of 44.9 hours of electrofishing in 2016. The two Des Moines River sites (Cliffland and confluence) accounted for 91.9% (171 individuals) of all Asian Carp captures, while the Skunk accounted for 6% (9 individuals), the Iowa River accounted for 1% (2 individuals), and Rock River accounted for 0.5% (1 individual). Silver Carp comprised 96% ($n=179$) of all Asian Carp captures in 2016, while Bighead Carp comprised 3% ($n=5$), and Grass Carp comprised 1% ($n=2$) of all captured individuals. Highest adult Asian Carp CPUE occurred at the confluence of the Des Moines River (Table 2). Confluence sites further upstream at the mouth of the Skunk, Iowa, Rock, and Wapsipinicon rivers had consecutively lower CPUE rates. Of the sites that were sampled twice (IAR-MTH and SKK-MTH), catch rates were slightly lower and the standard error increased in November compared to sampling done in September. The Des Moines River sites were the only sites to catch Bighead, Silver and Grass Carp. Only Silver Carp were caught at the Iowa, Skunk and Rock river confluences.

Des Moines River sites accounted for 94% of Silver Carp captures in total: 77% were captured at the Des Moines River confluence site ($n=137$) and 15% at the Cliffland site ($n=28$). The Skunk River accounted for 6% ($n=12$), the Iowa River accounted for 1% ($n=2$), and the Rock River accounted for 0.5% ($n=1$) of Asian Carp captures. Silver Carp ranged in size from 529 mm to 672 mm (mean = 619 mm; Figure 6) and 1.6 kg to 3.3 kg (mean = 2.3). Silver Carp size structure was smaller at Cliffland on the Des Moines River (DSM-CLF) compared to the Des Moines River confluence (Figure 6). The largest Silver Carp were captured at the the Skunk River confluence, ranging from 644 mm to 923 mm (mean = 829 mm; Figure 6) in length and 2.8 kg to 10.2 kg (mean = 6.9 kg) in weight. In addition to captures, few Asian Carp were observed leaping at sites outside of the Des Moines River that were unable to be captured due to jumping out of the range of the netters and the boat. A total of 41 Asian Carp jumps were observed at the Skunk River, four were observed at the Iowa River, and three were observed at the Rock River, but jumped out of range of the netters and boat.

Only four Bighead Carp were captured in 2016: three individuals were captured at the Des Moines River confluence and one individual was captured at Cliffland on the Des Moines River. Bighead Carp at the Des Moines River confluence ranged from 772 mm to 876 mm (mean = 832 mm) in length and 4.7 kg to 7.0 kg (mean = 5.9 kg) in weight. Only two Grass Carp were collected at Cliffland on the Des Moines River measuring 755 mm and 829 mm. Asian Carp otoliths collected during 2016 are currently being aged.

Environmental Condition Data

Chlorophyll *a* samples from the 2016 field season have all been processed (Figure 7). The confluence sites of each of the major tributaries had consistently higher chlorophyll *a* concentrations compared to upstream sites, downstream sites, and Pools 15 and 17 that lack a tributary. Similar chlorophyll *a* concentrations were generally documented among pools and in backwater, channel boarder, and thalweg habitats (Figure 7).

Products Resulting from this Project

Student theses

Camacho, C. A. 2016. Asian Carp reproductive ecology along the Upper Mississippi River invasion front. MS thesis, Iowa State University, Ames, Iowa.

Sullivan, C. J. 2016. Asian Carp population characteristics and dynamics in the Mississippi River watershed. MS thesis, Iowa State University, Ames, Iowa.

Matthews, A. In progress.

Tillotson, N. In progress.

Peer reviewed publications

Sullivan, C.J., C.A. Camacho, M.J. Weber, and C.L. Pierce. 2017. Intra-annual variability of Silver Carp populations in the Des Moines River, USA. *North American Journal of Fisheries Management* 37:836-849.

Sullivan, C.J., C.A. Camacho, M.J. Weber, C.L. Pierce, D.H. Wahl, and R. Columbo. In review. Factors regulating year-class strength of Silver Carp throughout North America. *Transactions of the American Fisheries Society*.

Camacho, C.A., C.J. Sullivan, M.J. Weber, and C.L. Pierce. In review. Suitability of an Upper Mississippi River Tributary for Asian Carp reproduction. *North American Journal of Fisheries Management*.

Sullivan, C.J., C.A. Camacho, M.J. Weber, C.L. Pierce, D.H. Wahl, and R. Columbo. In prep. Grass Carp occupancy and population characteristics in the Upper Mississippi River.

Camacho, C.A., C.J. Sullivan, M.J. Weber, and C.L. Pierce. In prep. Morphological identification of Asian Carp eggs using random forests machine learning classification.

Camacho, C.A., C.J. Sullivan, M.J. Weber, and C.L. Pierce. In prep. Asian Carp reproduction in the tributaries of the Upper Mississippi River.

Literature Cited

- Auer, N. A. (ed). 1982. Identification of larval fishes of the Great Lakes basin with emphasis on the Lake Michigan drainage. Great Lakes Fishery Commission, Ann Arbor, MI 48105. Special Pub 82-3:744 pp.
- Baerwaldt, K., A. Benson, and K. Irons. 2013. Asian Carp distribution in North America. Report to the Asian Carp Regional Coordinating Committee.
- Bruce, J. 1990. Bighead Carp (*Hypophthalmichthys nobilis*). Nonindigenous Aquatic Species Database, Gainesville, FL.
<http://nas.er.usgs.gov/queries/SpecimenViewer.aspx?SpecimenID=33085>, January 08, 2014. <http://nas.er.usgs.gov/queries/SpecimenViewer.aspx?SpecimenID=33085>.
- Burr, B. M., D. J. Eisenhour, K. M. Cook, C. A. Taylor, G. L. Seegert, R. W. Sauer, and E. R. Atwood. 1996. Nonnative fishes in Illinois waters: what do the records reveal. Transactions of the Illinois State Academy of Science 89(1):73–91.
- Camacho, C. A. 2016. Asian Carp reproductive ecology along the Upper Mississippi River invasion front. Masters thesis, Iowa State University.
- Chapman, D. C. 2006. Early development of four cyprinids native to the Yangtze River, China: U.S. Geological Survey Data Series 239.
- Chapman, D. C., and A. E. George. 2011. Developmental rate and behavior of early life stages of Bighead Carp and Silver Carp: U.S. Geological Survey Scientific Investigations Report 2011–5076.
- Crawley, M. J., H. Kornberg, J. H. Lawton, M. B. Usher, R. Southwood, R. J. O’Connor, and A. Gibbs. 1986. The population biology of invaders (and discussion). Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences 314(1167):711–731.
- DeGrandchamp, K. L., J. E. Garvey, and R. E. Colombo. 2008. Movement and habitat selection by invasive Asian Carps in a large river. Transactions of the American Fisheries Society 137(1):45–56.
- Deters, J. E., D. C. Chapman, and B. McElroy. 2013. Location and timing of Asian Carp spawning in the Lower Missouri River. Environmental Biology of Fishes 96(5):617–629.
- Freeze, M., and S. Henderson. 1982. Distribution and status of the Bighead Carp and Silver Carp in Arkansas. North American Journal of Fisheries Management 2(2):197–200.
- Garvey, J., B. Dugger, M. Whiles, S. Adams, M. Flinn, B. Burr, and R. Sheehan. 2003. Responses of fishes, waterbirds, invertebrates, vegetation, and water quality to environmental pool management: Mississippi River Pool 25. U.S. Army Corps of Engineers, St. Louis District, Final Report.
- Garvey, J. E., K. L. DeGrandchamp, and C. J. Williamson. 2006. Life history attributes of Asian Carps in the Upper Mississippi River system.
- Garvey, J., B. Ickes, and S. Zigler. 2010. Challenges in merging fisheries research and management: the Upper Mississippi River experience. Hydrobiologia 640(1):125–144.
- George, A. E., and D. C. Chapman. 2013. Aspects of embryonic and larval development in Bighead Carp (*Hypophthalmichthys nobilis*) and Silver Carp (*Hypophthalmichthys molitrix*). PLoS ONE 8(8):e73829.

- Gido, K. B., and J. H. Brown. 1999. Invasion of North American drainages by alien fish species. *Freshwater Biology* 42(2):387–399.
- Gozlan, R. E. 2008. Introduction of non-native freshwater fish: is it all bad? *Fish and Fisheries* 9(1):106–115.
- Hayer C. A., J. J. Breeggemann, R. A. Klumb, B. D. S Graeb, and K. N. Bertrand. 2014. Population characteristics of Bighead and Silver Carp on the northwestern front of their North American invasion. *Aquatic Invasions* 9: 289–303.
- Irons, K. 2012. Bighead Carp (*Hypophthalmichthys nobilis*). Nonindigenous Aquatic Species Database, Gainesville, FL.
<http://nas.er.usgs.gov/queries/SpecimenViewer.aspx?SpecimenID=543592>, January 08, 2014..
- Irons, K. S., G. G. Sass, M. A. McClelland, and T. M. O'Hara. 2011. Bigheaded Carp invasion of the La Grange Reach of the Illinois River: insights from the long term resource monitoring program. In *American Fisheries Society Symposium* (No. 74).
- Irons, K. S., S. A. DeLain, E. Gittinger, B. S. Ickes, C. S. Kolar, D. Ostendorf, E. N. Ratcliff, and A. J. Benson. 2009. Nonnative fishes in the Upper Mississippi River system: U.S. Geological Survey Scientific Investigations Report 2009–5176, 68 p.
<http://pubs.usgs.gov/sir/2009/5176/>.
- Isermann, D. A., W. L. McKibbin, and D. W. Willis. 2002. An analysis of methods for quantifying Crappie recruitment variability. *North American Journal of Fisheries Management* 22(4):1124–1135.
- Jennings, D. P. 1988. Bighead Carp (*Hypophthalmichthys nobilis*): a biological synopsis. U.S. Fish and Wildlife Service, Biological Report 88 (29).
- Kolar, C., D. Chapman, W. Courtenay, C. Housel, J. Williams, and D. P. Jennings. 2007. Bigheaded Carps: a biological synopsis and environmental risk assessment Special Publication 33. American Fisheries Society, Bethesda, Maryland.
- Krykhtin, M., and E. Gorbach. 1981. Reproductive ecology of the Grass Carp, *Ctenopharyngodon idella*, and the Silver Carp, *Hypophthalmichthys molitrix*, in the Amur Basin. *Journal of Ichthyology* 21(2):109–123.
- Larson, J. H., et al. 2017. Evidence of Asian Carp spawning upstream of a key choke point in the Mississippi River. *North American Journal of Fisheries Management* 37:903–917.
- Leuven, R. S. E. W., G. van der Velde, I. Baijens, J. Snijders, C. van der Zwart, H. J. R. Lenders, and A. bij de Vaate. 2009. The river Rhine: a global highway for dispersal of aquatic invasive species. *Biological Invasions* 11(9):1989–2008.
- Lohmeyer, A. M., and J. E. Garvey. 2009. Placing the North American invasion of Asian Carp in a spatially explicit context. *Biological Invasions* 11(4):905–916.
- Macisaac, H. J. 1996. Potential abiotic and biotic impacts of Zebra Mussels on the inland waters of North America. *American Zoologist* 36(3):287–299.
- Murphy, E. A., and P. R. Jackson. 2013. Hydraulic and water-quality data collection for the investigation of Great Lakes tributaries for Asian Carp spawning and egg-transport suitability: USGS Scientific Investigations Report 2013–5106. Page 30.
- Peters, L. M., M. A. Pegg, and U. G. Reinhardt. 2006. Movements of adult radio-tagged Bighead Carp in the Illinois River. *Transactions of the American Fisheries Society* 135(5):1205–1212.

- Pimentel, D., L. Lach, R. Zuniga, and D. Morrison. 2000. Environmental and economic costs of nonindigenous species in the United States. *BioScience* 50(1):53–65.
- Pimentel, D., R. Zuniga, and D. Morrison. 2005. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics* 52(3):273–288.
- Rahel, F. J. 2002. Homogenization of freshwater faunas. *Annual Review of Ecology and Systematics* 33:291–315.
- Rasmussen, J. L. 2002. The Cal-Sag and Chicago Sanitary and Ship Canal: a perspective on the spread and control of selected aquatic nuisance fish species. US Fish and Wildlife Service, Region 3.
- Sagoff, M. 2005. Do non-native species threaten the natural environment? *Journal of Agricultural and Environmental Ethics* 18(3):215–236.
- Sass, G. G., T. R. Cook, K. S. Irons, M. A. McClelland, N. N. Michaels, T. M. O’Hara, and M. R. Stroub. 2010. A mark-recapture population estimate for invasive Silver Carp (*Hypophthalmichthys molitrix*) in the La Grange Reach, Illinois River. *Biological Invasions* 12(3):433–436.
- Schrank, S. J., P. J. Braaten, and C. S. Guy. 2001. Spatiotemporal variation in density of larval Bighead Carp in the Lower Missouri River. *Transactions of the American Fisheries Society* 130(5):809–814.
- Southall, P. D., and W. A. Hubert. 1984. Habitat use by adult Paddlefish in the Upper Mississippi River. *Transactions of the American Fisheries Society* 113(2):125–131.
- Stohlgren, T. J., and J. L. Schnase. 2006. Risk analysis for biological hazards: what we need to know about invasive species. *Risk Analysis* 26(1):163–173.
- Sullivan, C. J. 2016. Asian Carp population characteristics and dynamics in the Mississippi River watershed. Masters thesis, Iowa State University, Ames, Iowa.
- Sullivan, C. J., C. A. Camacho, M. J. Weber, and C. L. Pierce. 2017. Intra-annual variability of Silver Carp populations in the Des Moines River, USA. *North American Journal of Fisheries Management* 37:836-849.
- Tweb, A., A. Ahmed, P. Baroi, and G. Mustafa. 1990. Studies on the development of the Silver Carp, *Hypophthalmichthys molitrix*. *Bangladesh Journal of Zoology* 18(2):139–145.
- U.S. Geological Survey. 2013. USGS water resources of the United States: boundary descriptions and names of regions, subregions, accounting units and cataloging units. http://water.usgs.gov/GIS/huc_name.html#Region07.
- United Press International. 2011. Bighead Carp (*Hypophthalmichthys nobilis*). Nonindigenous Aquatic Species Database, Gainesville, FL. <http://nas.er.usgs.gov/queries/SpecimenViewer.aspx?SpecimenID=275101>, January 08, 2014.
- United States Army Corps of Engineers (USACE). 2011. Waterborne commerce of the United States, Part 2—Waterways and Harbors Gulf Coast, Mississippi River System and Antilles. U.S. Army Corps of Engineers, IWR-WCUS-11-2.
- United States Army Corps of Engineers (USACE). 2014. The GLMRIS report: Great Lakes and Mississippi River interbasin study.

- Wanner, G., and R. Klumb. 2009. Asian Carp in the Missouri River: analysis from multiple Missouri River habitat and fisheries programs. National Invasive Species Council materials. Paper 10 National Invasive Species Council materials. Paper 10.
- Weber, M. J., and M. L. Brown. 2009. Effects of Common Carp on aquatic ecosystems 80 years after "Carp as a dominant": ecological insights for fisheries management. *Reviews in Fisheries Science* 17(4):524–537.
- Wilcox, D., E. Stefanik, D. Kelner, M. Cornish, D. Johnson, I. Hodgins, S. Zigler, and B. Johnson. 2004. Improving fish passage through navigation dams on the Upper Mississippi River System. US Army Corps of Engineers, ENV Report 54.
- Yi, B., Z. Yu, Z. Liang, S. Sujuan, Y. Xu, J. Chen, M. He, Y. Liu, Y. Hu, Z. Deng, S. Huang, J. Sun, R. Liu, and Y. Xiang. 1988. The distribution, natural conditions, and breeding production of the spawning ground on main stream of Yangtze River. In: Yi B, Yu Z, Liang Z (eds) *Gezhouba water control project and four famous fishes in the Yangtze River*. Hubei Science and Technology Press, Wuhan: 1–46.
- Zigler, S. J., M. R. Dewey, B. C. Knights, A. L. Runstrom, and M. T. Steingraeber. 2004. Hydrologic and hydraulic factors affecting passage of Paddlefish through dams in the Upper Mississippi River. *Transactions of the American Fisheries Society* 133(1):160–172.

Table 1. Eggs of 13 fish species captured in 2016 by date and river. River abbreviations present for a species on a sampling date denotes genetic confirmation of that species collected at that location on a sampling date. W= Wapsipinicon River, R= Rock River, I= Iowa River, S= Skunk River, D= Des Moines River, U= Upper Mississippi River.

	Common Name	Scientific Name	4/29/2016	5/10/2016	5/19/2016	5/29/2016	6/9/2016	6/18/2016	6/28/2016	7/8/2016	7/18/2016	7/29/2016	8/9/2016	8/18/2016	8/28/2016	9/9/2016
Asian Carp	Bighead Carp	<i>Hypophthalmichthys nobilis</i>				UIS										
	Silver Carp	<i>Hypophthalmichthys molitrix</i>				UIS										
	Grass Carp	<i>Ctenopharyngodon idella</i>				UIS						UI		U	U	
Other Species	Common Carp	<i>Cyprinus carpio</i>							UW							
	Emerald Shiner	<i>Notropis atherinoides</i>				UIS	UI	URIS	UIS	USD	UR	UD	UI	UI		
	Freshwater Drum	<i>Aplodinotus grunniens</i>	U	U	UR	URID	URISD	URIS	UIS	USD	USD	UI	UD	UD		
	Goldeye	<i>Hiodon alosoides</i>		D												
	River Shiner	<i>Notropis blennioides</i>							U	U			D			
	Sand Shiner	<i>Notropis stramineus</i>			UI		U	UDR			UID	UD	UI	UD		
	Shiner sp.	<i>Notropis sp.</i>								D						
	Shorthead Redhorse	<i>Moxostoma macrolepidotum</i>	D													
	Silvery Minnow	<i>Hybognathus nuchalis</i>				I										
	Speckled Chub	<i>Macrhybopsis aestivalis</i>				UIS	USD	U			UD		UD		D	

Table 2. Electrofishing catch per unit effort (CPUE, number per hour; mean \pm 1 SE) of all Asian Carp species (Bighead Carp, Silver Carp and Grass Carp) caught by location and date during 2016. Sites not sample during that month are marked with a (-).

Site	September	October	November
WAP-MTH	-	0	-
UMR-P15	-	0	-
ROC-MTH	-	0.35 \pm 0.35	-
UMR-P17	-	0	-
IAR-MTH	0.73	-	0.7 \pm 0.7
SKK-MTH	3.1 \pm 1.7	-	2.8 \pm 2.2
DSM-MTH	215.9 \pm 64.6	-	-
DSM-CLF	17.4	-	-




Figure 1. Approximate sampling (star) and dam (curved line) locations in rivers across southeastern Iowa. Sampling sites adjacent to tributaries of the Wapsipinicon, Rock, Iowa, Skunk and Des Moines rivers have sampling sites 1 km above and below the tributary within the Upper Mississippi River, in addition to a sampling site 1 km within the tributary.




Figure 2. Density (mean \pm 1 SE) of eggs (top) and age-0 fish (bottom) collected between April 29 and September 8, 2016.

*UMR-UPD, UMR-DND, and DSM-MTH were not sampled during this session.

**DSM-KQA was not sampled during this session.

2016




Figure 3. Densities (mean \pm 1 SE) of eggs (top) and age-0 fish (bottom) collected from each site during 2016. Site codes are expressed in the first three letters by the river they are sampled within: UMR=Upper Mississippi River: WAP = Wapsipinicon River: ROC=Rock River: IAR = Iowa River: SKK=Skunk River: DSM=Des Moines River. The second set of letter give a more descriptive location of the site. All UMR Sites contain either UP_, or DN_ corresponding to 1 km upstream and downstream 1 km of a major tributary. For example WAP-MTH = the Wapsipinicon River at the mouth river: UMR-UPW = Upper Mississippi River, upstream 1 km of the Wapsipinicon: UMR-DNW = Upper Mississippi River, downstream 1km of the Wapsipinicon River. The only tributary with two sites is the Des Moines River, where the second site above the mouth DSM-KQA is located at the town Keosauqua.

*One session from each of these siters were unable to be sampled.

Yr 2016




Figure 4. Egg densities (mean \pm 1 SE) captured from backwater, channel border, and thalweg habitats in the sites sampled during 2016. Below each habitat is the site code of the site sampled. UMR (Upper Mississippi River), MTH (mouth of the tributary) P15 (Pool 15 of the Mississippi River). See Figure 3 for a more complete site label descriptions.

*Not all of the Des Moines River sites were collected on 5/19/16 and 6/8/16 due to mechanical issues and low water levels.

Year 2016




Figure 5. Age-0 fish densities (mean \pm 1 SE) captured from backwater, channel border, and thalweg habitats in the sites sampled during 2016. Below each habitat is the site code of the site sampled. UMR (Upper Mississippi River), MTH (mouth of the tributary) P15 (Pool 15 of the Mississippi River). See Figure 3 for a more complete site label description.

*Not all of the Des Moines River sites were collected on 5/19/16 and 6/8/16 due to mechanical issues and low water levels.




Figure 6. Silver Carp length-frequency distributions and proportional size distribution (PSD; P-Preferred, M-Memorable, T-Trophy) indices of fish collected during 2016 from the Des Moines River at Cliffland (CLF) and at the Des Moines River mouth (MTH) in addition to Iowa River and Skunk River. One Silver Carp was captured at the Rock River mouth. No Asian Carp were captured in Pool 17, Pool 15, or the Wapsipinicon River confluence.




Figure 7. Chlorophyll *a* measurements (mean \pm 1 SE) of each site collected between April 29 and September 8, 2016.
 *Not all of the Des Moines River sites were collected on 5/19/16 and 6/8/16 due to mechanical issues and low water levels.